

## Antibacterial activity of tannins from the leaves of *Solanum trilobatum* Linn.

A. Doss, H. Mohammed Mubarak and R. Dhanabalan

Department of Microbiology, RVS College of Arts and Science, Coimbatore, Tamilnadu, India.

dossandro@gmail.com

**Abstract:** Compounds of pharmacological interest (tannins) were isolated from the plant species, *Solanum trilobatum* Linn and assayed against the bacteria, *Staphylococcus aureus*, *Streptococcus pyrogens*, *Salmonella typhi*, *Pseudomonas aeruginosa*, *Proteus vulgaris* and *Escherichia coli* using agar diffusion method. Tannins exhibited antibacterial activities against all the tested microorganisms. *S.aureus* was the most resistant to tannins isolated from the plant material followed by *Streptococcus pyrogens*, *Salmonella typhi*, *Escherichia coli*, *Proteus vulgari* and *Pseudomonas aeruginosa*. Minimum inhibitory concentration of the tannins ranged between 1.0 and 2.0 mg/ml while the minimum bactericidal concentration ranged between 1.5 and 2.0 mg/ml.

**Keywords:** *Solanum trilobatum*, tannins, methanol, bacterial pathogens, minimum inhibitory concentration, minimum bactericidal concentration

### Introduction

Tannins are polyphenols that are obtained from various parts of different plants (Gajendiran & Mahadevan, 1990). In addition to use in leather processing industries, tannins have shown potential antiviral (Lin *et al*, 2004), antibacterial (Akiyama *et al*, 2001; Funatogawa *et al*, 2004), and antiparasitic effects (Bhagavathi *et al*, 1999; Yang *et al*, 2000; Tanimura *et al*, 2005). In the past few years tannins have also been studied for their effects against cancer through different mechanisms.

*Solanum trilobatum* Linn. (Solanaceae-herbs) is an important medicinal plant. It contains rich amount of calcium, iron, phosphorus, carbohydrates, fat, crude fibre and minerals in the leaves (Jawahar *et al*, 2004). It is used to cure asthma, arrest blood vomiting, to reduce blood glucose level and bilious matter phlegmatic rheumatism and several kinds of leprosy. It is also antibacterial, antifungal, antimutagenic, antioxidant and antitumours (Purushothaman *et al*, 1969; Shahjahan *et al*, 2004; Shahjahan *et al*, 2005). This study was carried out to evaluate the antibacterial properties of tannins isolated from *S.trilobatum*.

### Materials and methods

#### Source of plant material

The leaves of the plant were collected from Western Ghats, Tamilnadu and the species was duly authenticated at Botanical Survey of India (Southern Circle), Coimbatore Tamilnadu.

#### Microorganisms

Clinical isolates of the bacteria viz., *Staphylococcus aureus*, *Streptococcus pyrogens*, *Salmonella typhi*, *Pseudomonas aeruginosa* *Proteus*

*vulgaris* and *Escherichia coli* used in this study were obtained from Microbial Diagnostic Laboratory, RVS Hospital, Coimbatore, Tamilnadu.

#### Preparation of plant extract

Methanolic extract of the leaf of the plant was prepared according to the method described by Okogun (2000).

#### Extraction of tannins

Sample of the powdered leaf (3 g) was boiled in 5 ml of distilled water for 3 min on a hot plate. The mixture was filtered while hot and the resulting filtrate was used to carry out ferric chloride test. (Trease & Evans, 1983). Sample of the filtrate (1.0 g) was transformed into a beaker and 10ml of distilled water was added. This was boiled for 5 min. Two drops of 5% ferric chloride (FeCl<sub>2</sub>) was then added. Production of greenish precipitate indicated the presence of tannins (Trease & Evans, 1983).

#### Antimicrobial test

The antibacterial test was performed by following agar disc diffusion method (Bauer & Kirby, 1996; Salie *et al*, 1996; Perez *et al*, 1990; Nair & Chanda, 2005) using Mueller Hinton Agar No. 2 medium for the assay. Microbial growth was determined by measuring the diameter of the zone of inhibition (SD ± Mean).

#### Determination of Minimum Inhibitory Concentration

Various concentrations of tannins from the plant, *Solanum trilobatum* ranging between 4.0 and 6.0 mg/ml were introduced into different test tubes; each tube was inoculated with an overnight culture of *Staphylococcus aureus*, *Proteus vulgaris*, *Streptococcus pyrogens*, *Escherichia coli*, *Salmonella typhi* and *Pseudomonas aeruginosa* diluted to give a final concentration of 10<sup>6</sup> cells per ml. The tubes were incubated at 37°C for 24 h. The least concentration of tannin that did not permit any visible growth of the inoculated test organism in broth culture was regarded as the minimum inhibitory concentration (MIC) in each case (Collins *et al*, 1995).

#### Determination of Minimum Bactericidal Concentration

After culturing the test organisms separately in nutrient broth containing various concentrations of the active ingredients, the broth was inoculated onto freshly prepared agar plates to assay for the bactericidal effect. The culture was incubated at 37°C for 24 h. The lowest concentration of alkaloid that does not yield any colony growth on the solid medium after the incubation period was regarded as minimum bactericidal concentration (MBC) (Alade & Irobi, 1995).



## Results and discussion

The presence of antibacterial substances in the higher plants is well established (Srinivasan *et al.*, 2001). Plants have provided a source of inspiration for

novel drug compounds as plants derived medicines have made significant contribution towards human health. Phytomedicine can be used for the treatment of diseases as is done in case of Unani and Ayurvedic system of medicines or it can be the base for the development of a medicine, a natural blueprint for the development of a drug (Jigna Parekh & Sumitra Chanda, 2007)

The sensitivity of seven pathogenic bacteria to the tannins of the leaves of *Solanum trilobatum* was tested and compared to that of antibacterial antibiotic Streptomycin. The results pointed out that these tannins possessed the highest antibacterial activity against the bacteria, *Staphylococcus aureus* and *Proteus vulgaris* at 2.5mg/ml concentration (Table.1), whereas they were moderately active against the remaining bacteria. Several plants which are rich in tannins have been shown to possess antimicrobial activities against a number of microorganisms. For example Banso and Adeyemo (2007) investigated the antibacterial activity of leaf extract *Dichrostachys cinerea* and reported that tannins, alkaloids and glycosides were detected. Amongst the gram-positive and gram-negative bacteria, gram-positive bacterial strains were more susceptible to the extracts when compared to gram negative bacteria. This may be attributed to the fact that these two groups differ by its cell wall component and its thickness (Yao *et al.*, 1995). The ability of tannin compounds to cause the bacterial colonies to disintegrate probably results from their interference with the bacterial cell wall; thereby inhibiting the microbial growth (Erasto *et al.*, 2004; Viljoen *et al.*, 2003).

The MIC of tannins isolated in this study against the test organisms ranged between 1.0 and 4.0 mg/ml while the MBC ranged between 1.5 and 4.5 mg/ml (Table 2). Antimicrobial agents with low activity against an organism had a high MIC while a highly active antimicrobial agent gave a low MIC. The antibacterial effectiveness with the increasing concentration of tannins observed in the present study is in agreement with the earlier investigations (Kurosaki & Nishi, 1983; Banso & Adeyemo, 2007). The results of the present study support the traditional use of the *Solanum trilobatum* as green medicine. It also suggests that the tannins isolated from the test plant possess remarkable

toxic activity against bacteria and may assume pharmacological importance.

Table 1. Antibacterial activity of tannin isolated from *S.trilobatum*.

Con. (mg/ml)	Mean diameter of zone of inhibition (mm ± SD)					
	<i>S. aureus</i>	<i>P.vulgaris</i>	<i>P. aeruginosa</i>	<i>S.typhi</i>	<i>S. pyrogens</i>	<i>E.coli</i>
0.5	8.3 ± 0.28	NA	NA	NA	NA	NA
1	8.6 ± 0.28	7.8 ± 0.280	NA	8.5 ± 0.5	9.6 ± 0.28	NA
1.5	10.5 ± 0.5	7.83 ± 0.15	8.0 ± 0.5	9.13 ± 0.32	9.6 ± 0.52	8.8 ± 0.26
2	11.5 ± 0.5	8.83 ± 0.28	7.83 ± 0.76	10.5 ± 0.5	10.5 ± 0.5	8.8 ± 0.35
2.5	13.5 ± 0.5	9.66 ± 0.28	8.73 ± 0.25	10.8 ± 0.28	11.5 ± 0.5	10.5 ± 0.05
Strepto-mycin	20	22	17	16	18	21

NA - No Activity

Table 2. Minimum inhibitory concentration (MIC) and Minimum bactericidal concentration (MBC) of tannins isolated from *Solanum trilobatum*.

Clinical isolates	MIC (mg/ml)	MBC (mg/ml)
<i>Staphylococcus aureus</i>	1	1.5
<i>Proteus vulgaris</i>	4	4.5
<i>Pseudomonas aeruginosa</i>	4	4.5
<i>Salmonella typhi</i>	2	2.0
<i>Streptococcus pyrogens</i>	2	2.5
<i>Escherichia coli</i>	4	4.5

## References

1. Akiyama H, Kazuyasu F, Yamasaki O, Oono T and Iwatsuki K (2001) Antibacterial action of several tannins against *Staphylococcus aureus*. *J. Antimicrobial Chemotherapy*, 48 (48), 487-491.
2. Alade PI and Irobi ON (1995) Antifungal activities of crude leaf extract of *Acalypha wilesiana*. *J. Ethnopharmacol.* 39, 171-174.
3. Banso A and Adeyemo SO (2007) Evaluation of antibacterial properties of tannins isolated from *Dichrostachys cinerea*. *Afr. J. Biotechnol.* 6 (15), 1785-1787.

