

Effect of Bay Laurel (*Laurus nobilis L.*) Leaf Powder Dietary Supplementation on Dressing Percent, Carcass Traits, Carcass Cuts and Some Internal Organs of Quail

Mohammed Ali Makki Al Rubae*

Department of Animal Production, College of Agriculture, Wasit University Iraq; poultman76@yahoo.com

Abstract

Background/Objective: Bay laurel (*Laurus nobilis L.*) leaf can be used in poultry nutrition to increase gastric fluids to stop digestive disorders; it has antifungal, antibacterial potentials. The leaves had many effective substances including Apigenin, Luteolin, Kaempferol, Myricetin, cinole, eugenol, terpenol, Glycosides, Quercitin, linalool, Alkaloids, methyl eugenol and lactonol. **Methods:** This study was conducted to measure the Effect of bay laurel (*Laurus nobilis L.*) leaf powder dietary supplementation on dressing percents, carcass trait, carcass cuts and some internal organs of meat type quail by 0, 1, 2, 3% to the diets fed to 180 seven days old meat type quail randomly distributed to four treatments with three replicates for each treatment (15 bird /replicate). **Findings:** the bay laurel (*Laurus nobilis L.*) leaf powder dietary supplementation significantly by increased live body weight, carcass weight, dressing percent, and percentage weights of breast, drumstick, and thigh cuts. Adding bay laurel leaf powder improves taste, tenderness, and total acceptance of breast meat. **Improvements:** all experimental diets (of bay laurel (*Laurus nobilis L.*) leaf powder dietary supplementation) resulted in good dressing percents, carcass trait, carcass cuts in quail meat, indicating that this feed supplement should be used on industrial scale.

Keywords: Bay Laurel Leaf, Carcass, Cuts, Dressing Percents, Quail

1. Introduction

The new international trend to decrease or ban of antibiotics and chemicals in poultry industry was the motive for researchers to find new sources of growth promoters, antioxidants, remedies and anti inflammation materials¹. Thus herbs or photogenic were introduced as active components of the diet who have the ability to boost bird's immunity and performance².

Bay laurel (*Laurus nobilis L.*) leaf can be used in poultry nutrition to increase gastric fluids to stop digestive disorders³. It has antifungal, antibacterial potentials^{4,5}. The leaves had many effective substances including Apigenin, Luteolin, Kaempferol, Myricetin, cinole, eugenol, terpenol⁶⁻⁸. Glycosides, Quercitin, lin-

alool, Alkaloids, Methyl Eugenol, lactonol (Go`mez-Glycosides).

There were a few studies about adding Bay laurel (*Laurus nobilis L.*) leaf oil broiler's diets⁹⁻¹⁰ and broiler breeder¹¹ in quail¹² has reported that supplementation of bay laurel leaves (2 or 4 g/kg feed) didn't effect on performance or egg quality parameters of quails, the study recorded that supplemented the diet with bay laurel leaves is able to change the biochemical traits of quail egg yolk. Very few scientific experiments done to study the effect of bay laurel leaves on meat type quails and carcass composition. The aim of this current experiment is to study the effect of bay laurel (*Laurus nobilis L.*) leaf powder dietary supplementation on dressing percents, carcass trait, carcass cuts and some internal organs of meat type quail.

*Author for correspondence

2. Materials and Methods

Dry bay laurel (*Laurus nobilis L.*) leaf was purchased from a local market in Baghdad, Iraq and grounded to powder for further use. A total of 180 seven day old quails were raised according to social and behavioral research ethics committee. Chicks were weighed individually and randomly allocated to 4 dietary treatments. Birds had free access to feed and water. Experiment was carried out in a completely randomized design with 4 treatments; each had 3 replicates with 15 birds/replicate. Diets were formulated according to the requirements of NRC¹³ (Table 1). The experimental groups were as follows: Group I (control) fed a basal diet, Group II fed the basal diet + 1% bay laurel (*Laurus nobilis L.*) leaf powder, Group III fed the basal diet + 2% bay laurel (*Laurus nobilis L.*) leaf powder, Group IV fed the basal diet + 3% bay laurel (*Laurus nobilis L.*) leaf powder. Continuous lighting program (23 hours lightning: 1hour darkness) was provided. At 49 days of age five birds from each treatment group were slaughtered for measuring carcass weight and dressing percentage, carcass cuts, femur and sternum bones length, body length, intestine length, body index, thigh index, heart and lever and gizzard percentage weight, breast diameter, thigh diameter¹⁴. Sensory meat quality parameters, evaluation was conducted by 25 trained panelists. Breast meat cuts from each treatment were tested as described by¹⁵. Using the table of degrees (Table 2) data analyzed with ANalysis Of VARIances (ANOVA) followed by¹⁶ Differences between treatments were analyzed using Statistical Analysis System¹⁷.

3. Results and Discussion

The results presented in Table 3 shows a significant increase in live body weight by supplementing 2% bay

Table 1. Dietary^{*} composition of experimental diet

Ingredients (g/kg)	%
Maize	48
Soybean meal	38
Protein concentrate**	10
Vegetable oil	3
Limestone	0.7
Salt	0.3
Total	100
Protein %	25.8
ME (Kilocalorie/kg)	2932.74
Met + sys.	0.7
Ca	2.54
Available p.	0.35

*NRC¹³, **Protein concentrates (50% protein, 2800 kilocalorie, 12% fat, 25% ash, 2.5% ca, 2.9% p)

laurel (*Laurus nobilis L.*) leaf powder which had the highest live body weight by 159.78 gm followed by the rest of the experimental treatments, while control treatment recorded the lowest live body weight. The same differences were noticed for carcass weight and dressing percent. Control birds had significantly the heaviest gizzard and liver weight by 3.06 and 2.91 respectively no differences were reported for heart percentage weight and intestine length.

No significant differences were noticed for body degree of fullness, barest degree of fullness, thigh diameter, sternum length, and thigh bone length (Table 4) control treatment birds had the shortest ($p < 0.5$) body length, and breast diameter 3% bay laurel leaf powder treatment was longer ($p < 0.5$) for leg bone length.

In Table 5 we can find many differences in carcass cuts percentage weights bay laurel leaf powder supple-

Table 2. Degrees of breast meat sensory test

Degree	Odor	Color	Juiciness	Tenderness	Odor	Total acceptance
1.	Very good	Very good	Very good	Very good	Very good	Very good
2.	Good	Good	Good	Good	Good	Good
3.	Medium	Medium	Medium	Medium	Medium	Medium
4.	Bad	Bad	Bad	Bad	Bad	Bad
5.	Very bad	Very bad	Very bad	Very bad	Very bad	Very bad

Table 3. The effect of bay laurel (*Laurus nobilis L.*) leaf powder dietary supplementation on live body weight cm, carcass weight cm, dressing percents, and gizzard, heart, liver percentage weight and intestine length of quail

TRT	Live body weight cm	Carcass weight cm	Dressing percent %	Gizzard percentage weight	Heart percentage weight	Liver percentage weight	Intestine length cm
1.	143.26±10.11c	95.39±9.35b	66.30±2.05b	3.06±0.65a	1.47±0.20a	2.91±0.21a	54.00±2.00a
2.	159.78±7.61a	109.51±6.26a	68.47±1.03a	2.65±0.30b	1.40±0.08a	2.78±0.26b	51.16±2.20a
3.	149.06±5.71b	97.42±5.61b	65.18±1.85b	2.78±0.13b	1.40±0.08a	2.72±0.84b	52.50±1.75a
4.	152.87±7.99ab	102.90±6.40ab	67.25±0.98ab	2.64±0.28b	1.45±0.09a	2.74±0.94b	52.60±1.80a

*The different letters in the same column indicate a significant difference

Table 4. The effect of bay laurel (*Laurus nobilis L.*) leaf powder dietary supplementation on body length, body degree of fullness, breast degree of fullness, Breast diameter, thigh diameter, leg bone length, sternum bone length, thigh bone length of quail

TRT	Breast%	Drumstick%	Thigh%	Wings%	Back%	Neck%
1.	34.14±0.57b	8.44±0.30b	14.24±1.92b	8.48±0.34b	26.92±0.59a	5.37±0.91a
2.	34.82±0.76ab	9.65±0.31a	15.73±1.29a	9.49±0.11a	28.80±0.72a	5.42±0.29a
3.	34.70±0.37ab	9.18±0.34ab	15.07±0.61a	9.76±0.53a	27.15±0.62a	5.47±1.08a
4.	36.94±0.92a	9.06±0.20ab	14.84±0.50ab	9.66±0.21a	26.61±2.20a	5.53±0.69a

*The different letters in the same column indicate a significant difference

Table 5. The effect of bay laurel (*Laurus nobilis L.*) leaf powder dietary supplementation on breast, thigh diameter, leg bone length, wings, back, and neck percentage weight of the carcass of quail

TRT	Body length cm	Body degree of fullness	Breast degree of fullness	Breast diameter cm	Thigh diameter cm	leg bone length cm	Sternum length cm	Thigh bone length cm
1.	7.64±0.30b	12.40±0.89a	1.57±0.07a	11.03±0.25b	4.92±0.13a	3.82±0.02b	3.36±0.04a	4.62±0.07a
2..	8.26±0.10a	12.17±0.60a	1.62±0.18a	12.26±0.45a	5.22±0.09a	3.96±0.06ab	3.31±0.24a	4.85±0.09a
3.	8.71±0.34a	12.95±0.66a	1.53±0.03a	12.85±0.62a	5.34±0.18a	3.94±0.05ab	3.41±0.07a	4.84±0.03a
4.	8.16±0.30a	12.59±0.66a	1.53±0.08a	12.49±0.58a	5.13±0.36a	4.06±0.06a	3.41±0.08a	4.79±0.07a

*The different letters in the same column indicate a significant difference

mentation treatments surpassed the control treatment in percentage weights of breast, drumstick, thigh, wings no differences were found for percentage weights of back and neck. Sensory traits of breast meat results (Table 6) shown that 3% bay laurel leaf powder treatment had the best ($p < 0.5$) taste, tenderness, and total acceptance than the other treatments. Most of the supplementation treatments were significantly higher than control treatments for appearance, odor, color and juiciness.

Japanese quail birds have a small condensed body compacted with meat¹⁸, they have a 60–80% dressing percent which is higher than broiler's. The results of the current study agrees with those of¹⁹ who found a significant increase in live body weight by supplementing essential oils to quail diets, and with the results of²⁰ who fed essen-

tial oils of bay laurel (*Laurus nobilis L.*) contains linlool, cineole, methyl eugenol which are present in bay laurel (*Laurus nobilis L.*) leaf powder^{21,22}

There is high positive correlation between live body weights and dressing percent²³ so by the increase in live body weight we had an increase in dressing percent, this agrees with the results of this current study. Same trend of differences in live body weight was noticed for dressing percent. The improvement of carcass yield and carcass cuts is attributed live body weight improvement^{24,25}.

The dressing percent results in our results agrees with²⁶ results in improving dressing percent and carcass cuts by adding herbal mix. It also agrees with the findings of²⁷ who added herbal mixtures to quail diets, he recorded an increased metabolic rate and bone and muscle develop-

Table 6. The effect of bay laurel (*Laurus nobilis L.*) leaf powder dietary supplementation on breast meat appearance, taste, odor, color, juiciness, tenderness, total acceptance of quail

TRT	Appear-ance	Taste	Odor	Color	Juiciness	Tender-ness	Total acceptance
1.	3.67 b	3.83b	3.50b	3.70b	3.60b	3.80b	3.87b
2.	3.87a	3.87b	3.80ab	3.77b	3.80ab	3.80b	3.83b
3.	3.77ab	3.87b	3.83ab	4.00a	4.00a	.87b3	3.87b
4.	3.73ab	4.00a	4.07a	3.90ab	4.00a	3.93a	3.97a

*The different letters in the same column indicate a significant difference

ment which lead to increased dressing percent by herbal mixture supplementation .

Eugenol in bay laurel (*Laurus nobilis L.*) leaf powder enhanced the growth of beneficial bacteria²⁸ like lactobacillus and depress the harmful bacteria such as *E. coli* in the intestine, this action can improve general health and nutrient availability and performance in poultry²⁹ bay laurel (*Laurus nobilis L.*) leaf powder had a high content of minerals that can motivate many enzymatic systems in the liver and consolidate metabolism³⁰. This explains the decreased liver and gizzard weights by bay laurel (*Laurus nobilis L.*) leaf powder supplementation, these results didn't agree with those of³¹ who didn't notice any differences in liver and gizzard weights when feeding bay laurel (*Laurus nobilis L.*) oil to quail birds.

The higher bone length result in the current study is an index to higher live body weight and bay laurel (*Laurus nobilis L.*) leaf powder content³¹. Long bones means long skeletal muscles especially in t and thigh cuts³² bay laurel (*Laurus nobilis L.*) leaf has been used as food condiment, it's used to give cooked meat flavor and tenderness. Our present results of sensory breast meat traits agrees with those of^{33,34}, both fed herbal mix to broilers and found improvement in sensory traits of meat, this might be due increased water holding capacity of the breast meat of birds fed with bay laurel (*Laurus nobilis L.*) leaf powder in which the polyphenols that can ameliorate meat quality^{35,36}. Also the addition of herbs can inhibit poultry meat spoilage by bacteria and give positive sensory perception³⁷.

4. Conclusion

All The dietary supplementation levels (1, 2, and 3%) of bay laurel (*Laurus nobilis L.*) leaf powder had a positive effect on dressing percents, carcass trait, carcass cuts of quail birds.

5. References

1. Tipu MA, Pasha TN, ALL Z. Comparative efficacy of alinomyacin sodium and Neem fruit (*Azadirachta indica*) as feed additive Anticoccidials in broilers, International Journal of Poultry Science. 2006; 1(4):91–93.
2. Esonu BO, Emenalom OO, Udedibie ABI, Anyanwu GA, Madu U, Inyang AO. Evaluation of Neem (*Azadirachta indica*) leaf meal on performance, carcass characteristics and egg quality of laying hens, International Journal of Agricultural Development. 2005; 6(1):208–12.
3. Matsuda H, Shimoda H, Ninomiya K, Yoshikawa M. Inhibition mechanism of costunolide, a sesquiterpene lactone isolated from *Laurus nobilis*, on blood-ethanol elevation in rats: Involvement of inhibition of gastric emptying and increase in gastric juice secretion, Alcohol. 2002; 37(2):121–27. <https://doi.org/10.1093/alcalc/37.2.121>.
4. Kumar S, Singh J, Sharma A. Bay leaves. In: Peter, K.V (Ed.) Handbook of Herbs and Spices. Woodhead Publishing Limited, Cambridge, UK; 2001. p. 52–61. <https://doi.org/10.1533/9781855736450.52>. PMID: PMC2955933
5. Muhammad A, Dangoggo SM, Tsafe AI, Itodo AU, Atiku FA. Proximate, minerals and Anti-nutritional factors of *Gardenia aqualla* (*Gaudendutse*) fruit pulp. Pakistan Journal of Nutrition. 2011, 10 (6), pp. 577-581. <https://doi.org/10.3923/pjn.2011.577.581>.
6. Gomez-Coronado D, Barbas C. Optimized and validated HPLC method for R- and G-tocopherol measurement in *Laurus nobilis* leaves new data on tocopherol content, Journal of Agricultural and Food Chemistry. 2003; 51(18):5196–201. <https://doi.org/10.1021/jf030143f>. PMID: 12926858.
7. Conforti F, Starri G, Uzunov D, Menichini F. Comparative chemical composition and antioxidant activities of wild and cultivated *Laurus nobilis L.* leaves and *Foeniculum vulgare* subsp – (*Ucria*) Coutinho seeds, Biological and Pharmaceutical Bulletin. 2006; 29(10):2056–64. <https://doi.org/10.1248/bpb.29.2056>. PMID: 17015951.
8. Yasar U, Ibrahim IO, Ibrahim EY, Iihan D, Goksel D. Determination of some heavy metals and mineral nutri-

- ents of bay tree (*Laurus nobilis* L.) in Bartın city, Turkey, *Pakistan Journal of Botany*. 2012; 44:81–89.
9. Osman M, Yakout HM, Mot-Awe HF, Ezz-ElArab WF. Productive, Physiological, Immunological and Economical effect of supplementing natural feed additives to broiler diets, *Egyptian Poultry Science Journal*. 2010; 30(1):25–53.
 10. Ozek K. Effects of dietary herbal essential oil mixture and/or mannan-oligosaccharide supplementation on laying performance, some serum biochemical markers and humoral immunity in laying hens exposed to heat, *Revue de Médecine Vétérinaire*. 2012; 163(3):153–59.
 11. Traesel CK. Serum biochemical Profile and performance of broiler chickens fed diet containing essential oils and pepper, *Comparative Clinical Pathology*. 2011; 20(5):453–60. <https://doi.org/10.1007/s00580-010-1018-1>.
 12. Karaalp M, Mahfuz E, Nusret G, Metin S, Musa Y, Mehmet O. Bay laurel (*Laurus nobilis* L.) in Gapanese Quails feeding 1, performance and egg quality parameters, *Journal of Animal and Veterinary Advances*. 2011; 10(14):1883–89. <https://doi.org/10.3923/javaa.2011.1883.1889>.
 13. National Research Council (NRC). *Nutrient Requirement of Poultry*. 9th Revised Edition. National Academy Press. Washington, DC; 1994. p. 1–176.
 14. USDA. U.S. Department of Agriculture, Agricultural Research Service. USDA National Nutrient Database for Standard Reference; Release 26, 2013.
 15. Jayasena DD, Jung S, Kim, HJ, Bae, YS, Yong HI, Lee JH, Kim JG, Jo CR. Comparison of quality traits of meat from Korean native chickens and broilers used in two different traditional Korean cuisines, *Asian-Australian Journal of Animal Sciences*. 2013; 26(7):1038–46. <https://doi.org/10.5713/ajas.2012.12684>. PMID: 25049883, PMCID: PMC4093503.
 16. Duncan DB. Multiple ranges and multiple F-test, *Biometrics*. 1955; 11(1):1–42. <https://doi.org/10.2307/3001478>.
 17. SAS. *User's Guide. Statistical. Version 9. 1sd Edition. Statistical Analysis System. SAS. Institute Inc. Cary. N.C. USA; 2012.*
 18. Rogerio CT. Quail meat undiscovered alternative, *World Poultry*. 2009; 25(2):12–14.
 19. Biricik H, Yesilbag D, Gezen SS, Bulbul T. Effects of dietary myrtle oil (*Myrtus communis* L.) supplementation on growth performance, meat oxidative stability, meat quality and erythrocyte parameters in quails, *Revue de Médecine Vétérinaire*. 2012; 163(3):131–38.
 20. Alciçek A, Bozkurt M, Cabuk M. The effect of an essential oil combination derived from selected herbs growing wild in Turkey on broiler performance, *South African Journal of Animal Science*. 2003; 33(2):89–94. <https://doi.org/10.4314/sajas.v33i2.3761>.
 21. Ozek T, Demirci B, Baser KHC. Chemical composition of Turkish Myrtle oil, *Journal of Essential Oil Research*. 2000; 12(5):541–44. <https://doi.org/10.1080/10412905.2000.9712154>.
 22. Kucukyilmaz K, Catli AU, Cinar M. Etlikpiliçyemlerinees ansiyelyağkarışımıilavesininbüyüme Performansı, Karkas Randımanıve Bazıç organ Ağırıkları Üzerine Etkileri, *Kafkas Üniversitesi Veteriner Fakültesi Dergisi*. 2012; 18(2):291–96.
 23. Zangana BSR, Al-Mashhadani SM. Effect of adding different levels from marjoram (*Origanum vulgare*) powder to the diet in some of carcass measurement of broiler, *Alanbar Journal of Agricultural Science*. 2017; 15(1):426–34.
 24. Yesilbag D, Gezen SS, Biricik H, Bulbul T. Effect of a rosemary and oregano volatile oil mixture on performance, lipid oxidation of meat and haematological parameters in pharaoh quails, *British Poultry Science*. 2012; 53(1):89–97. <https://doi.org/10.1080/00071668.2012.654763>. PMID: 22404809.
 25. Bulbul T, Vural O, Aziz B. Use of sage (*Salvia triloba* L.) and laurel (*Laurus nobilis* L.) oils in quail diets, *Eurasian Journal of Veterinary Sciences*. 2015; 31(2):95–101. <https://doi.org/10.15312/EurasianJVetSci.2015210080>.
 26. Simsek UG, Ciftci M, Dalkilic B, Ertas GT. The effects of dietary antibiotic and anise oil supplementation on body weight, carcass characteristics and organoleptic analysis of meat in broilers, *Revue de Médecine Vétérinaire*. 2007; 158(10):514–18.
 27. AL-Khateeb FS. Effect of dietary inclusion of ginger, garlic and black bean seeds on some productive traits of Japanese Quail, *Jordan Journal of Agricultural Sciences*. 2017; 13(2):465–76.
 28. Agostini PS, Sola-Oriol D, Nofrarias M, Barroeta AC, Gasa J, Manzanilla EG. Role of in-feed clove supplementation on growth performance, intestinal microbiology, and morphology in broiler chicken, *Livestock Science*. 2012; 147(1-3):113–18. <https://doi.org/10.1016/j.livsci.2012.04.010>.
 29. Ordóñez G, Llopis N, Penalver P. Efficacy of Eugenol against a *Salmonella enterica* serovar enteritidis experimental infection in commercial layers in production, *Journal of Applied Poultry Research*. 2008; 17(3):376–82. <https://doi.org/10.3382/japr.2007-00109>.
 30. Weber GM, Michalczuk M, Huyghebaert G, Juin H, Kwakernaak C, Gracia MI. Effects of a blend of essential oil compounds and benzoic acid on performance of broiler chickens as revealed by a meta-analysis of growth trials in various locations, *The Journal of Poultry Science*. 2012; 91(11):2820–28. <https://doi.org/10.3382/ps.2012-02243>. PMID: 23091138.

31. Gwaad RA, Gwaad RA. Effect of aqueous extract of *Laurus nobilis* L. leaves on some physiological parameters and testis tissue in male white rabbit *lepus articus*, *Journal of Kerbala University*. 2017; 11(3):31–41.
32. Al-Hajo NN. The Effect of Uropygiectomy on Some Quality Characteristics and Sensory Evaluation of Broiler Meat in Different Age. Ph.D. thesis, University of Baghdad, Department of Animal Resources; 2010.
33. Toghyani M, Gheisari A, Ghalmkari G, Mohammadrezaei M. Growth performance, serum biochemistry and blood hematology of broiler chicks fed different levels of black seed (*Nigella sativa*) and peppermint (*Mentha piperita*), *Livestock Science*. 2010; 129(1-3):173–78. <https://doi.org/10.1016/j.livsci.2010.01.021>.
34. Al-Sultan SI. The effect of *Curcuma longa* (turmeric) on overall performance of broiler chickens, *International Journal of Poultry Science*. 2003; 2(5):351–53. <https://doi.org/10.3923/ijps.2003.351.353>.
35. Liu XF, Xia YN, Fang Y. Effect of metal ions on the interaction between bovine serum albumin and berberine chloride extracted from a traditional Chinese herb *Coptischinensis* Franch, *Journal of Inorganic Biochemistry*. 2005; 99(7):1449–57. <https://doi.org/10.1016/j.jinorg-bio.2005.02.025>. PMID: 15908003.
36. Bostoglou NA, Florou-Paneri P, Christaki E, Fletouris DJ, Spais AB. Effect of dietary oregano essential oil on performance of chickens and on iron- induced lipid oxidation of breast, thigh and abdominal fat tissues, *British Poultry Science*. 2002; 43(2):223–30. <https://doi.org/10.1080/00071660120121436>. PMID: 12047086.
37. Samant S, Crandall PG, O'Bryan C, Lingbeck JM, Martin EM, Seo HS. Sensory impact of chemical and natural antimicrobials on poultry products: A review, *Poultry Science*. 2015; 94(7):1699–710. <https://doi.org/10.3382/ps/pev134>. PMID: 26015590.